PhytoKeys

PhytoKeys 265: 209–221 (2025)
DOI: 10.3897/phytokeys.265.167921

Research Article

Indosasa fimbriligulata (Poaceae, Bambusoideae), a new bamboo species from Guangxi, China

Zheng-Yang Niu[1,2], Lin Bai[1,2], Yi-Hua Tong[1,2], Nian-He Xia[1,2]

1 *State Key Laboratory of Plant Diversity and Specialty Crops & Guangdong Provincial Key Laboratory of Digital Botanical Garden, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, 510650, China*

2 *South China National Botanical Garden, Chinese Academy of Sciences, Guangzhou, 510650, China*

Corresponding author: Yi-Hua Tong (yh-tong@scbg.ac.cn)

Abstract

Based on morphological and molecular phylogenetic evidence, we confirmed that some bamboo collections from Guangxi, China, represent a new species of *Indosasa*. The new species, named *I. fimbriligulata*, is morphologically similar to *I. glabrata* but differs by having internodes that are white strigose when young, and hollow without or with a little clastic pith, culm buds triangular, culm leaf sheaths abaxially scattered with tufted brown setae, culm leaf auricles well developed, and culm leaf ligule apex long fimbriate. A detailed description, as well as two color plates of this new species, is also provided.

Key words: Arundinarieae, morphology, phylogeny, taxonomy, three-branched bamboos

Introduction

Indosasa McClure was established by F. A. McClure (1940), with *I. crassiflora* designated as its generic type. It belongs to the subtribe Arundinariinae of the tribe Arundinarieae (Poaceae, Bambusoideae) (Zhang et al. 2020) and contains 17 species mainly distributed in China (Fujian, Hunan, Guangdong, Guangxi, Guizhou, Yunnan), Vietnam, and Laos (Zhu and Zhao 1996; Zhu and Stapleton 2006; Vorontsova et al. 2016; Niu et al. 2021, 2022, 2025a). It is diagnosed by leptomorph (monopodial) rhizomes, diffuse culms, strongly raised culm supra-nodal ridges, pseudospikelets, six stamens, and three stigmas per floret (McClure 1940; Chao and Chu 1983; Zhu and Zhao 1996; Xue et al. 2003; Zhu and Stapleton 2006; Yi et al. 2008; Xia and Lin 2009; Xia et al. 2016; Shi et al. 2021). However, some recent phylogenomic studies suggested that *Indosasa* sensu lato (s.l.) is non-monophyletic (Guo et al. 2021; Niu 2025). *Indosasa* sensu stricto (s.s.) is defined by the possession of strongly raised and asymmetrically swollen branch supra-nodal ridges, usually 4–9 foliage leaves per ultimate branch, and the traits mentioned above (Niu 2025). The newly defined *Indosasa* s.s. has only eight species, and its distribution is limited to Guangxi, Yunnan, and South Guizhou of China, North Vietnam, and North Laos (Niu 2025).

During two field investigations in Fangchenggang City, Guangxi, we encountered a running bamboo that does not fit the description of any known species.

OPEN ACCESS

Academic editor: Wei Lim Goh
Received: 6 August 2025
Accepted: 15 September 2025
Published: 4 November 2025

Citation: Niu Z-Y, Bai L, Tong Y-H, Xia N-H (2025) *Indosasa fimbriligulata* (Poaceae, Bambusoideae), a new bamboo species from Guangxi, China. PhytoKeys 265: 209–221. https://doi.org/10.3897/phytokeys.265.167921

Copyright: © Zheng-Yang Niu et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

However, some traits, such as strongly raised culm supra-nodal ridges, strongly raised and asymmetrically swollen branch supra-nodal ridges, and the branch complement with three branches per mid-culm node, match well with those diagnostic of *Indosasa* s.s. After comparison with all the previously recorded species of *Indosasa* in national and local floras (Chao and Chu 1983; Zhu and Zhao 1996; Xue et al. 2003; Zhu and Stapleton 2006; Yi et al. 2008; Xia and Lin 2009; Xia et al. 2016; Shi et al. 2021), as well as the monograph of *Indosasa* (Niu 2025), we found that this species is most similar to *I. glabrata* C.D.Chu & C.S.Chao. In order to clarify the taxonomic identity of this bamboo, we conducted phylogenetic and morphological studies. Finally, we confirmed that this bamboo is new to science based on morphological and molecular evidence.

Materials and methods

Morphological study

Morphological comparisons between this new species and morphologically similar species were conducted based on living plants in the field, specimens, protologues, and descriptions from floras (Chao and Chu 1983; Zhu and Zhao 1996; Zhu and Stapleton 2006; Xia et al. 2016; Niu 2025). Some detailed characters, such as indumentum, were observed with a stereomicroscope (Mshot MZ101, Guangzhou Micro-shot Technology Co., Ltd., Guangzhou, China). Measurements were taken using a ruler or micrometer. Terminology follows McClure (1940), Li et al. (2006), and Beentje (2016).

Taxon sampling, DNA extraction, and sequencing

To ascertain the phylogenetic position of this new species, we conducted phylogenetic analyses based on nuclear single-copy orthologous genes (SOGs). According to the most recent phylogenetic framework of three-branched bamboos (Niu et al. 2025b), we selected 22 samples representing 21 bamboo species belonging to eight genera. Specifically, the ingroup from the tribe Arundinarieae contained six species of *Indosasa*, three species of *Acidosasa* B.M.Yang, three species of *Oligostachyum* Z.P.Wang & G.H.Ye, three species of *Pseudosasa* Makino ex Nakai, two species of *Pleioblastus* Nakai, two species of *Sinobambusa* Makino ex Nakai, and one species of *Indocalamus* Nakai. *Bambusa vulgaris* Nees from the tribe Bambuseae was chosen as the outgroup. Sample numbers and their corresponding voucher information are listed in Suppl. material 1: table S1.

For DNA extractions, young leaves were collected in the field and dried with silica gel. Total genomic DNA was extracted from the dried leaves using the Rapid Plus DNA Lib Prep Kit for Illumina (RK2008) (ABclonal, Woburn, USA) and sheared (1 μg DNA per sample) using a Covaris M220 ultrasonicator (Covaris, Woburn, USA). PCR products were purified using the AMPure XP system (Beverly, USA). Library quality was assessed on the Agilent 5400 system (Agilent, USA), and DNA quantification was performed by quantitative polymerase chain reaction (qPCR). The qualified paired-end library was pooled using the NEBNext® Ultra™ DNA Library Prep Kit and sequenced on the NovaSeq 6000

platform with the PE150 strategy. After filtration of adapters and low-quality reads using Fastp software v. 0.23.2 (Chen et al. 2018), we obtained at least 40 Gb of deep genome skimming data.

Target nuclear single-copy orthologous genes capture

For nuclear gene recovery, we used the protein-coding sequences of six previously published bamboo genomes – *Dendrocalamus latiflorus* Munro (Zheng et al. 2022), *Phyllostachys edulis* (Carrière) J.Houz. (Zhao et al. 2018), *Bonia amplexicaulis* (L.C.Chia, H.L.Fung & Y.L.Yang) N.H.Xia, *Guadua angustifolia* Kunth, *Olyra latifolia* L., and *Raddia guianensis* (Brongn.) Hitchc. (Guo et al. 2019) – to identify 737 common nuclear single-copy orthologous genes (SOGs) using Orthofinder v. 2.5.4 (Emms and Kelly 2019). We recovered putative SOGs using HybPiper v. 2.0.1 (Johnson et al. 2016). In this pipeline, we employed BWA v. 0.7.17 (Li and Durbin 2009) to map filtered reads to each specified SOG. We then de novo assembled reads and mapped them to each SOG into contigs using SPAdes v. 3.15.0 (Bankevich et al. 2012). We aligned the assembled contigs to the reference SOG dataset and used the Python script 'retrieve_sequences.py' to recover 737 putative orthologs for each sample. However, because our samples are polyploid (Guo et al. 2019), some so-called SOGs may have multiple copies. We therefore used the Python script 'paralog_retriever.py' to detect and remove potential paralogs. After this step, we retained 479 SOGs.

Alignment and phylogenetic inference

We aligned the 479 SOGs using MAFFT v. 7.505 (Katoh and Standley 2013) in Geneious v. 9.1.4 (Kearse et al. 2012). We trimmed each single-gene matrix using trimAl v. 1.4 (Salvador et al. 2009) with default settings. We then removed nuclear genes with lengths shorter than 300 bp or with >50% missing data. The final nuclear dataset used for phylogenetic analyses included 465 SOGs. All SOG alignments can be found in Suppl. material 1.

We used a multi-species coalescent-based method for nuclear gene-based phylogenetic inference. We first inferred individual ML trees using RAxML v. 8.2.12 (Stamatakis 2014) for each nuclear gene and estimated branch support using bootstrapping analysis (Felsenstein 1985) with 500 replicates, all under the GTRGAMMAI model. To improve coalescent analysis accuracy (Zhang et al. 2018; Cai et al. 2021), we collapsed branches with support values below 30% using Newick Utilities v. 1.6.0 (Junier and Zdobnov 2010). We used the resulting collapsed individual gene trees and corresponding bootstrapped trees to infer the species tree based on the multi-locus bootstrapping (MLB) method in ASTRAL-III (Seo 2008; Zhang et al. 2018). We then created 500 coalescent bootstrap replicates, each consisting of 465 gene trees (one tree per gene, randomly sampled from bootstrapping trees without replacement) for quartet analysis. The final MLB species tree was inferred from the 500 resulting "species" trees. MLB branch bootstrap values (BS) were calculated based on the frequency with which branches appeared across the 500 resulting "species" trees using the greedy consensus method (Soto Gomez et al. 2019). We considered BS >70% as strong support.

Results

Our nuclear SOG-based MLB species tree strongly supported that the new species is a member of *Indosasa* s.s. and is closely related to *I. glabrata* (BS = 100) (Fig. 1).

Morphological comparison revealed that the new species is morphologically similar to *I. glabrata* in having small culms that are 2–4 m high and 0.5–2 cm in diam., narrow and long triangular papery culm leaf sheaths, and linear-lanceolate culm leaf blades, whereas the other *Indosasa* species are usually characterized by relatively large culms, leathery culm leaf sheaths, and broad lanceolate to ovate culm leaf blades. The new species can be easily distinguished from *I. glabrata* by having triangular (vs. oblate) culm buds, white strigose (vs. glabrous) young culm internodes, without pith or with slightly clastic pith (vs. lamellate pith), abaxial surface of culm leaf sheath with scattered clumps of brown setae and base of culm leaf sheath with dense brown setae (vs. glabrous culm leaf sheath), well-developed (vs. undeveloped) culm leaf auricles and oral setae, and culm leaf ligule with long fimbriae (vs. glabrous) (Table 1; Figs 2–4).



Figure 1. The multi-locus bootstrapping ASTRAL species tree shows the phylogenetic position of *Indosasa fimbriligulata*, which is reconciled by coalescence of 465 single-copy orthologous nuclear gene trees after collapsing branches with support values <30%. BS support values below 70% are represented by a slash.

Table 1. Morphological comparison of *Indosasa fimbriligulata* and *I. glabrata*.

Characters	*I. fimbriligulata*	*I. glabrata*
Young culm		
Primary buds	Triangular	Oblate
Surface of internodes	White strigose	Glabrous
Cavity pith	Nearly absent	Lamellate
Culm leaf		
Abaxial surface of sheaths	Scattered with tufted brown setae	Glabrous
Base of abaxial surface of sheaths	Densely brown setose	Glabrous
Auricles	Small, ca. 0.5 × 1 mm	Undeveloped
Oral setae	Developed, 2–10 mm long	Undeveloped
Apex of ligules	Fimbriate	Glabrous
Blade shape	Linear to linear-lanceolate	Narrow lanceolate to lanceolate

Discussion

Although recent phylogenetic studies based on plastid and nuclear sequences confirmed that *Indosasa* s.l. is non-monophyletic (Ma et al. 2017; Guo et al. 2021; Niu 2025), this new species is undoubtedly a member of *Indosasa* s.s. based on both morphological and molecular evidence (Figs 1–4; Table 1). Not only do its morphological characters match well with those of *Indosasa* s.s., but the phylogenetic analysis also showed that it and *I. glabrata* form a monophyletic clade with the type species of *Indosasa*, viz., *I. crassiflora*. Further comparison with morphologically similar species confirmed that it is a distinct new species, which is described and illustrated below.

Taxonomic treatment

Indosasa fimbriligulata **N.H.Xia, Z.Y.Niu & Y.H.Tong, sp. nov.**
urn:lsid:ipni.org:names:77371342-1
Figs 2–4

Type. CHINA • Guangxi Zhuang Autonomous Region: Fangchenggang City, Naliang Town, Gaolin Village; 21°37'48"N, 107°40'57"E; alt. 391 m; 17 April 2025; *L. Bai & H. Y. Li BLSC-25041501* (holotype: IBSC!).

Diagnosis. *Indosasa fimbriligulata* resembles *I. glabrata* but differs from the latter in having white strigose internodes when young, hollow internodes without pith or with a little clastic pith, abaxial surface of culm leaf sheaths scattered with tufted brown setae, well-developed culm leaf auricles, and culm leaf ligule apex with long fimbriae.

Description. Rhizomes leptomorph. Culms diffuse, erect, sometimes flexuose, 2–4 m tall and 0.5–1.5 cm in diameter; internodes flattened on branching side, with two longitudinal ridges and three grooves above branching points along the branching side, 30–50 cm long, initially green to dark green, yellow-green when aged, white strigose when young, glabrescent when old, white powdery only at infranodal regions; walls 3–4 mm thick, cavity hollow without pith or with slightly

Figure 2. *Indosasa fimbriligulata* N.H.Xia, Z.Y.Niu & Y.H.Tong. **A.** Habit; **B.** Old culm; **C.** Culm bud; **D.** Branch complement with three branches at culm node; **E.** Portion of culm leaf showing leaf blade; **F.** Portion of culm leaf showing sheath base; **G.** Culm pith. Scale bars: 5 cm (**G**).



Figure 3. *Indosasa fimbriligulata* N.H.Xia, Z.Y.Niu & Y.H.Tong. **A.** Branch complement, showing strongly raised and asymmetrically swollen branch supra-nodal ridges; **B.** Foliage leafy branchlets; **C.** Ultimate foliage leafy branchlet; **D.** Portion of foliage leafy branchlet showing auricles and oral setae; **E.** Portion of culm leaf, showing culm leaf blade; **F.** Culm leaf ligule with long fimbriae; **G.** Culm leaf sheath base. Scale bars: 5 cm (**A**); 10 cm (**C**); 5 mm (**F**); 1 cm (**D, E, G**).



Figure 4. *Indosasa fimbriligulata* N.H.Xia, Z.Y.Niu & Y.H.Tong. **A.** Culm leaf sheath; **B.** Portion of culm, showing leaf blade, node, and internode; **C.** Culm node, showing culm bud and sheath scar; **D.** Branch complement with three branches at culm node, showing strongly raised and asymmetrically swollen branch supra-nodal ridges; **E.** Ultimate foliage leafy branchlet. Scale bars: 2 cm (**A–E**).

clastic pith; supra-nodal ridges strongly raised, sheath scars prominent, with a corky ring and densely brown setose when young, glabrescent when old, with persistent remains of sheath base. Branch supra-nodal ridges strongly raised and asymmetrically swollen. Primary buds solitary, triangular, yellowish green, not sunken into culm, margin densely ciliate. Branch complement with three branches at each culm node. Culm leaf sheaths initially green with small purple dots, more so on the marginal part, turning to straw when old, tardily deciduous, narrow and long triangular, papery, abaxially scattered with tufted brown setae, densely brown setose at base, margins densely ciliate; auricles small, ovate to triangular, ca. 1 × 0.5 mm; oral setae well-developed, many, straight or slightly tortuous, 2–10 mm long; ligules arcuate, ca. 2 mm tall, abaxially puberulent, apex long fimbriate, fimbriae many, straight or slightly tortuous, 3–7 mm long; blades reflexed, linear to linear-lanceolate, 0.8–3 × 0.1–0.3 cm, 3/50–6/50 as long as sheath, apex acuminate, base narrowed, adaxially sparsely brown setae, abaxially glabrous. Foliage leaves 4–7 per ultimate branchlet; sheaths 3.5–6.5 cm long, glabrous, longitudinal ribs conspicuous when dry; auricles well-developed, ovate to triangular, 3–5 × 1–3 mm; oral setae many, straight or slightly tortuous, 3–20 mm long; ligules short, truncate, 0.5–1 mm tall, abaxially pubescent; blades lanceolate, papery, 9.5–24 × 1.5–3.8 cm, apex acuminate, base cuneate to widely cuneate, glabrous on both sides, one margin serrulate, other margin entire, secondary veins 5–8 pairs, transverse veinlets conspicuous. Inflorescence unknown.

Phenology. New shoots from April to June.

Distribution and habitat. This new species has been found only in its type locality so far. It usually occurs on mountain slopes near roadsides and streams.

Etymology. The specific epithet refers to the fimbriate culm leaf ligule of this species, which is rarely seen in other *Indosasa* species. Its Chinese name is given as 流苏大节竹 [liú sū dà jié zhú].

Additional specimens examined. China • Guangxi: Fangchenggang City, Naliang Town, Gaolin Village; 21°37'34"N, 107°41'8"E; alt. 443 m; 15 June 2025; *Z. Y. Niu*, *NZY297* (IBSC).

Local usage. Old culms are used for fences.

Acknowledgments

We are grateful to Ms. Hong-Yan Li (South China Botanical Garden, Chinese Academy of Sciences) for her assistance during the fieldwork and to Mr. Ding-Han Cui (South China Botanical Garden, Chinese Academy of Sciences) for the line drawing.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Use of AI

No use of AI was reported.

Funding

This study was supported by the National Natural Science Foundation of China (Grant No. 32270227), the Chinese Academy of Sciences (Biotaxonomic Scientist Post Grant No. CAS-TAX-24-049), the Basic Research Projects Independently Deployed by South China Botanical Garden, Chinese Academy of Sciences (Grant No. JCYJXM-202507), and the National Wild Plant Germplasm Resource Center.

Author contributions

Zheng-Yang Niu: writing original manuscript, field investigation, data analysis. Lin Bai: field investigation. Yi-Hua Tong: funding support, data acquisition, reviewing, and editing the manuscript. Nian-He Xia: supervision, funding support, reviewing, and editing the manuscript.

Author ORCIDs

Zheng-Yang Niu https://orcid.org/0000-0003-0281-1504
Yi-Hua Tong https://orcid.org/0000-0002-5034-005X
Nian-He Xia https://orcid.org/0000-0001-9852-7393

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA (2012) SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. Journal of Computational Biology : A Journal of Computational Molecular Cell Biology 19(5): 455–477. https://doi.org/10.1089/cmb.2012.0021

Beentje H (2016) The Kew Plant Glossary: An Illustrated Dictionary of Plant Terms (Second Edition). Royal Botanic Gardens Kew Press, Richmond, Surrey, 1–184.

Cai LM, Xi ZX, Lemmon EM, Lemmon AR, Mast A, Buddenhagen E, Liu L, Davis CC (2021) The Perfect Storm: Gene Tree Estimation Error, Incomplete Lineage Sorting, and Ancient Gene Flow Explain the Most Recalcitrant Ancient Angiosperm Clade, Malpighiales. Systematic Biology 70(3): 491–507. https://doi.org/10.1093/sysbio/syaa083

Chao CS, Chu CD (1983) A study on the bamboo genus *Indosasa* of China. Zhiwu Fenlei Xuebao 21(1): 60–75.

Chen SF, Zhou YQ, Chen YR, Gu J (2018) fastp: An ultra-fast all-in-one FASTQ preprocessor. Bioinformatics (Oxford, England) 34(17): 884–890. https://doi.org/10.1093/bioinformatics/bty560

Emms DM, Kelly S (2019) OrthoFinder: Phylogenetic orthology inference for comparative genomics. Genome Biology 20(1): 238. https://doi.org/10.1186/s13059-019-1832-y

Felsenstein J (1985) Confidence limits on phylogenies: An approach using the bootstrap. Evolution; International Journal of Organic Evolution 39(4): 783–791. https://doi.org/10.2307/2408678

Guo ZH, Ma PF, Yang GQ, Hu JY, Liu YL, Xia EH, Zhong MC, Zhao L, Sun GL, Xu YX, Zhao YJ, Zhang YC, Zhang YX, Zhang XM, Zhou MY, Guo Y, Guo C, Liu JX, Ye XY, Chen YM, Li DZ (2019) Genome sequences provide insights into the reticulate origin and

unique traits of woody bamboos. Molecular Plant 12(10): 1353–1365. https://doi.org/10.1016/j.molp.2019.05.009

Guo C, Ma PF, Yang GQ, Ye XY, Guo Y, Liu JX, Liu YL, Eaton DAR, Guo ZH, Li DZ (2021) Parallel ddRAD and genome skimming analyses reveal a radiative and reticulate evolutionary history of the temperate bamboos. Systematic Biology 70(4): 756–773. https://doi.org/10.1093/sysbio/syaa076

Johnson MG, Gardner EM, Liu Y, Medina R, Goffinet B, Shaw AJ, Zerega NJC, Wickett NJ (2016) HybPiper: Extracting coding sequence and introns for phylogenetics from high-throughput sequencing reads using target enrichment. Applications in Plant Sciences 4(7): 1600016. https://doi.org/10.3732/apps.1600016

Junier T, Zdobnov EM (2010) The Newick utilities: High-throughput phylogenetic tree processing in the UNIX shell. Bioinformatics (Oxford, England) 26(13): 1669–1670. https://doi.org/10.1093/bioinformatics/btq243

Katoh K, Standley DM (2013) MAFFT: multiple sequence alignment software version 7: improvements in performance and usability. Molecular Biology and Evolution 30(4): 772–780. https://doi.org/10.1093/molbev/mst010

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A (2012) Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics (Oxford, England) 28(12): 1647–1649. https://doi.org/10.1093/bioinformatics/bts199

Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics (Oxford, England) 25(14): 1754–1760. https://doi.org/10.1093/bioinformatics/btp324

Li DZ, Wang ZP, Zhu ZD, Xia NH, Jia LZ, Guo ZH, Yang GY, Stapleton CMA (2006) Bambuseae (Poaceae). In: Wu ZY, Raven PH, Hong DY (Eds) Flora of China (Vol. 22) Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 7–180.

Ma PF, Vorontsova MS, Nanjarisoa OP, Razanatsoa J, Guo ZH, Haevermans T, Li DZ (2017) Negative correlation between rates of molecular evolution and flowering cycles in temperate woody bamboos revealed by plastid phylogenomics. BMC Plant Biology 17(1): 260. https://doi.org/10.1186/s12870-017-1199-8

McClure FA (1940) New genera and species of Bambusaceae from eastern Asia. Lingnan University Science Bulletin 9: 1–67.

Niu ZY (2025) Phylogenomic study and taxonomic revision of *Indosasa* (Poaceae: Bambusoideae). PhD Dissertation, University of Chinese Academy of Sciences 1–375.

Niu ZY, Tong YH, Xia NH (2021) *Pleioblastus triangulata* (Poaceae: Bambusoideae), a new combination for *Indosasa triangulata* based on morphological and molecular evidence. Phytotaxa 527(4): 275–283. https://doi.org/10.11646/phytotaxa.527.4.5

Niu ZY, Cai ZY, Liao CL, Xia NH (2022) *Chimonobambusa sangzhiensis* (Poaceae, Bambusoideae), a new combination supported by morphological and molecular evidence. PhytoKeys 195: 127–141. https://doi.org/10.3897/phytokeys.195.83004

Niu ZY, Cai ZY, Yin J, Tong YH, Xia NH (2025a) Morphological and molecular evidence suggests that *Indosasa gigantea* and *Acidosasa glauca* (Poaceae, Bambusoideae, Arundinarieae) are conspecific. PhytoKeys 255: 171–187. https://doi.org/10.3897/phytokeys.255.143020

Niu ZY, Zhang ZX, Cai ZY, Tong YH, Zeng SJ, Graham SW, Xia NH (2025b) Phylogenomic and morphological evidence supports the reinstatement of the bamboo genus *Clavinodum* from *Oligostachyum* (Poaceae: Bambusoideae). Molecular Phylogenetics and Evolution 207: 108327. https://doi.org/10.1016/j.ympev.2025.108327

Salvador C, José MS, Toni G (2009) trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics (Oxford, England) 25(15): 1972–1973. https://doi.org/10.1093/bioinformatics/btp348

Seo TK (2008) Calculating bootstrap probabilities of phylogeny using multilocus sequence data. Molecular Biology and Evolution 25(5): 960–971. https://doi.org/10.1093/molbev/msn043

Shi JY, Zhang YS, Zhou DQ, Ma LS, Yao J (2021) Illustrated Flora of Bambusoideae in China (Vol. 1). Science Press & Springer, Beijing, 1–637.

Soto Gomez M, Pokorny ML, Kantar MB, Forest F, Leitch IJ, Gravendeel B, Wilkin P, Graham SW, Viruel J (2019) A customized nuclear target enrichment approach for developing a phylogenomic baseline for *Dioscorea* yams (Dioscoreaceae). Applications in Plant Sciences 7(6): e11254. https://doi.org/10.1002/aps3.11254

Stamatakis A (2014) RaxML version 8: A tool for phylogenetic analysis and post analysis of large phylogenies. Bioinformatics (Oxford, England) 30(9): 1312–1313. https://doi.org/10.1093/bioinformatics/btu033

Vorontsova MS, Clark LG, Dransfeld J, Govaerts R, Baker WJ (2016) World checklist of bamboos and rattans. INBAR Technical Report, No. 37. International Network of Bamboo & Rattan, Beijing, 1–454.

Xia NH, Lin RS (2009) Poaceae. I. Bambusoidese. In: Wu TL (Ed.) Flora of Guangdong (Vol. 9). Guangdong Provincial Publishing Group, Guangdong Science and Technology Press, Guangzhou, 224–325.

Xia NH, Dai QH, Tang SC (2016) Poaceae. I. Bambusoidese. In: Li SG (Ed.) Flora of Guangxi (Vol. 5). Guangxi Science & Technology Publishing House, Nanning, 739–823.

Xue JR, Yang YM, Li DZ (2003) *Indosasa*. In: Sun BX, Li DZ, Xue JR (Eds) Flora Yunnanica (Vol. 9). Science Press, Beijing, 173–178.

Yi TP, Shi JY, Ma LS, Wang HT, Yang L (2008) Iconographia Bambusoidearum Sinicarum. Science Press, Beijing, 1–766.

Zhang C, Rabiee M, Sayyari E, Mirarab S (2018) Astral III: Polynomial time species tree reconstruction from partially resolved gene trees. BMC Bioinformatics 19(S6): 153. https://doi.org/10.1186/s12859-018-2129-y

Zhang YX, Guo C, Li DZ (2020) A new subtribal classifcation of Arundinarieae (Poaceae, Bambusoideae) with the description of a new genus. Plant Diversity 42(3): 127–134. https://doi.org/10.1016/j.pld.2020.03.004

Zhao HS, Gao ZM, Wang L, Wang JL, Wang SB, Fei BH, Chen CH, Shi CC, Liu XC, Zhang HL, Lou YF, Chen LF, Sun HY, Zhou XQ, Wang SN, Zhang C, Xu H, Li LC, Yang YH, Wei YL, Yang W, Gao Q, Yang HM, Zhao SC, Jiang ZH (2018) Chromosome-level reference genome and alternative splicing atlas of moso bamboo (*Phyllostachys edulis*). GigaScience 7(10): giy115. https://doi.org/10.1093/gigascience/giy115

Zheng YS, Yang DM, Rong JD, Chen LG, Zhu Q, He TY, Chen LY, Ye J, Fan LL, Gao YB, Zhang HX, Gu LF (2022) Allele-aware chromosome-scale assembly of the allopolyploid genome of hexaploid Ma bamboo (*Dendrocalamus latiflorus* Munro). Journal of Integrative Plant Biology 64(3): 649–670. https://doi.org/10.1111/jipb.13217

Zhu ZD, Stapleton CMA (2006) *Indosasa*. In: Wu ZY, Raven P (Eds) Flora of China (Vol. 22). Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 143–147.

Zhu ZD, Zhao QS (1996) *Indosasa*. In: Geng BJ, Wang ZP (Eds) Flora Reipublicae Popularis Sinicae (Vol. 9). Science Press, Beijing, 204–224.

Supplementary material 1

465 single-copy orthologous gene alignments used for the reconstruction of MLB ASTRAL species tree and one supplementary table

Author: Zheng-Yang Niu

Data type: zip

Explanation note: Here are 465 single-copy orthologous gene alignments used for the reconstruction of multi-locus bootstrapping ASTRAL species tree, and their corresponding specimen voucher information used in our study.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/phytokeys.265.167921.suppl1